

iTrack[™] to be Featured at the 2022 American Glaucoma Society (AGS) Annual Meeting

Fremont, California, 3 March 2022 – Nova Eye Medical Limited, a medical technology company committed to advanced ophthalmic treatment technologies and devices, is pleased to announce that the Company's proprietary iTrack[™] canaloplasty technology will be featured at the American Glaucoma Society (AGS) Annual Meeting, being held on March 3-6, 2022, in Nashville, Tennessee.

A summary of the data to be presented during the official scientific program of the AGS is outlined below:

Poster Presentation:

Mahmoud A Khaimi, MD

Canaloplasty Standalone Versus Combined with Cataract Surgery- 36-month Outcomes in 1000+ eyes

Robert Noecker, MD, MBA

12-Month Evaluation of Endothelial Cell Density and Loss Following Canaloplasty

Poster presentation details can be viewed in the AGS 2022 program book (refer to pages 70 and 71) which can be accessed via the AGS website: <u>https://www.americanglaucomasociety.net/ags-events/annual-meeting</u>



ABOUT NOVA EYE MEDICAL

Nova Eye Medical Limited is a medical technology company that develops, manufactures and sells a portfolio of proprietary ophthalmic treatment technologies and devices. Used by eye surgeons in more than 100 countries globally, these technologies include iTrack[™], a consumable surgical device for the treatment of glaucoma. The Company also manufactures and sells the proprietary Molteno3[®] glaucoma drainage device for the treatment of severe or complex glaucoma. With its sales headquarters based in Fremont, California, Nova Eye Medical is supported by sales offices in Adelaide, Australia and Berlin, Germany, and a global network of more than 50 distribution partners. Manufacturing facilities are located in Fremont, California and Dunedin, New Zealand.

For additional information about Nova Eye Medical and its technologies, please visit: <u>www.nova-eye.com</u>

ABOUT CANALOPLASTY

First introduced in 2008, canaloplasty is a surgical treatment for glaucoma that targets the main sites of outflow resistance in the conventional outflow pathway: the trabecular meshwork, Schlemm's canal, and the distal collector channels. Based on the same principles as angioplasty, a flexible microcatheter is cannulated 360 degrees around Schlemm's canal during the procedure to manually break and remove blockages. Next, viscoelastic fluid is injected into Schlemm's canal as the microcatheter is withdrawn to dilate the distal outflow system and to improve the function of the trabecular meshwork.

Canaloplasty is typically performed using either of the following two surgical techniques:

• Performed via an ab-interno surgical technique, canaloplasty is a highly effective treatment option for cases of mild-moderate glaucoma. It typically reduces intraocular pressure (IOP) to the low teens. It has also



been observed to reduce patient dependence on medications. The abinterno surgical technique is an implant-free, tissue-sparing procedure that preserves future treatment options.

Performed via an ab-externo surgical technique, canaloplasty is a highly
effective treatment option for patients with severe glaucoma that
overcomes the risks and discomfort associated with traditional glaucoma
surgery. With over 100,000 procedures performed to date, clinical
studies show that canaloplasty has an excellent safety profile, with
minimal post-operative follow-up, fast recovery time, and infrequent
intra-operative and postoperative complications.

ABOUT THE iTRACK™ PORTFOLIO

Nova Eye Medical (formerly iScience Interventional) pioneered the canaloplasty market with the launch of the world's first canaloplasty device, $iTrack^{TM}$, in 2008. Since then, more than 100,000 canaloplasty procedures have been performed with the $iTrack^{TM}$ device, cementing its role in the treatment of glaucoma both as a standalone procedure and in combination with cataract surgery.

The iTrackTM canaloplasty microcatheter has been cleared for the indication of fluid infusion and aspiration during surgery, and for catheterization and viscodilation of Schlemm's canal to reduce intraocular pressure in adult patients with openangle glaucoma. The iTrackTM canaloplasty microcatheter is currently not 510(k) cleared for use with the ab-interno technique in the United States,

For additional information about iTrack[™], including safety information, please visit: <u>www.glaucoma-iTrack.com</u>